

Supplementary Information

Association between *PPAR-γ2* Pro12Ala genotype and insulin resistance is modified by circulating lipids in Mexican children

Carolina Stryjecki¹, Jesus Peralta-Romero², Akram Alyass¹, Roberto Karam-Araujo³, Fernando Suarez², Jaime Gomez-Zamudio², Ana Burguete-Garcia⁴, Miguel Cruz^{2,*}, David Meyre^{1,5}

¹Department of Clinical Epidemiology and Biostatistics, McMaster University, Hamilton, ON, Canada

²Medical Research Unit in Biochemistry, Hospital de Especialidades, Centro Médico Nacional Siglo XXI del Instituto Mexicano del Seguro Social, Mexico City, Mexico

³Health Promotion Division, Instituto Mexicano del Seguro Social. Mexico City, Mexico

⁴Centro de investigación sobre enfermedades infecciosas. Instituto Nacional de Salud Pública. Cuernavaca, Morelos, Mexico

⁵Department of Pathology and Molecular Medicine, McMaster University, Hamilton, ON, Canada


Supplementary Table S1: Power calculation for the main effect of *PPAR-γ2* rs1801282 on BMI.

MAF	Beta	Sample Size (unadjusted) ^a	Sample Size (adjusted) ^b	Interaction Sample Size (unadjusted) ^a	Interaction Sample Size (adjusted) ^b
0.14	0.50	2296	4495	118	230
	0.60	1596	3119	79	155
	0.70	1169	2290	56	109
	0.80	894	1751	41	80
	0.90	706	1382	31	60

Abbreviations: MAF, minor allele frequency

^a power calculation unadjusted for multiple testing (2 sided p-value = 0.05, 80% power)

^b power calculation adjusted for multiple testing (2 sided p-value = 2.08×10^{-3} , 80% power)



Supplementary Figure S1: Histograms illustrating raw distribution (panel A) and corrected distributions following inverse normal transformations (panel B) of variables of interest.